

Phytoremediation



AECOM phytoremediation specialists consist of a multidisciplinary team of ecologists, botanists, engineers and subject matter experts versed in plant assisted remediation.

Areas of Expertise

- Site suitability screening and plant selection
- Cost value analysis
- Mechanistic approaches
- Plant uptake studies and potted plant growth studies
- Emerging contaminant uptake and transformation
- Conceptual approach and design
- Field and performance monitoring services
- Pilot- and full-scale implementation


More Information:

Barry J. Harding, CPG, Director Nature Based Solutions, Remediation West | Barry.Harding@aecom.com
 Clara Foutch, Project Manager & Ecologist, Remediation West | Clara.Foutch@aecom.com

Overview

Phytoremediation is the applied use of plants to transform, degrade or remove contaminants from environmental media, and is a nature-based remedial solution. The use of phytoremediation systems for the treatment of contaminants in groundwater, process wastewaters, surface water, and soil has become increasingly popular as a result of their effectiveness, relatively low capital and operation & maintenance costs, and increased regulatory acceptance. AECOM has been at the forefront of using cost-effective phytoremediation technologies in the United States and several other countries. Phytoremediation has been proven as a long-term, cost-effective remedial technique for controlling off-site migration of groundwater through phyto-hydraulics, and aerobically-degrading amenable contaminants such as petroleum, benzene, toluene, ethylbenzene and xylene (BTEX) and polycyclic aromatic hydrocarbons compounds. In addition, phyto-technologies can also be applied to other wastestream treatments including metals (phytoextraction and phytosequestration), energetic compounds, and emerging compounds such as 1,4-dioxane. Per and polyfluoroalkyl substances (PFAS) can also be extracted into foliar tissues of plants through phytoextraction mechanisms.

Phytoremediation systems are gaining acceptance by the regulatory community and include the added benefits of providing site stabilization, visual screening, and habitat restoration. Such systems are unobtrusive, typically do not interfere with other remediation systems, and can be a supplement or cornerstone of a site's long-term remedial closure and site restoration plan. Whether the application be remediation of contamination in soil and groundwater, hydraulic control for reduction of off-site groundwater migration, or the polishing of a wastewater stream prior to reintroduction to the environment, AECOM strives to design effective and cost-efficient systems using proven, phytoremediation technologies.



Groundwater phytoremediation decision tree

Determining Cost-Effective Applications

Plants take 1-3 years to become established.

Depends on the required thickness of capture-zone and the permeability of the aquifer.


Tree water-use declines in winter. Plumes can be controlled year round by adjusting shape and size of the tree stand.

Special engineered practices can train tree roots down to 25-ft.

Level of tolerance depends on contaminant types, concentrations, and plant species. Greenhouse studies may be necessary to assess an appropriate plant type.

LEGEND

- (A) BTEX**
BTEX compounds are degraded by microbes residing in the plant root-zone (rhizosphere).
- (B) TCE/Dioxane**
TCE and 1,4-dioxane can be taken up, translocated and exit plant shoots (phytovolatilization). This may be followed by UV-degradation in the atmosphere.
- (C) TNT/RDX**
TNT and RDX; following plant uptake, these compounds are metabolized and detoxified by plant enzymes.
- (D) Some Metals**
Some metals can safely accumulate in the woody tissue of trees.
- (E) Some Pesticides**
Depending on conditions, for sites contaminated with chlordane.



Phytoremediation *continued*



Natural remediation using salt-tolerant trees



Natural remediation using indigenous grasses

Our approach

Phytoremediation technologies. AECOM uses various Molecular Biological Tools (MBTs) to evaluate the presence, behavior and performance of bacteria and archaea residing in both the soil rhizosphere, aquifer, and plant tissues. The approach is to establish environmental baselines regarding the microbial community and use a weight of evidence evaluation in assessing the performance of plant-assisted remediation in a modern omics-driven era.

Enhanced rhizodegradation (plant-assisted bioremediation). Enhanced rhizodegradation, or plant-assisted bioremediation is an enhanced biodegradation method for treating organic chemical pollutants by root-dwelling bacteria and fungi living under the influence of select plant species. This biodegradation method is facilitated by a “rhizosphere effect” where plant roots improve soil quality through bioturbation and increased aeration, increased porosity, deposit carbon-containing compounds (rhizodeposition) into the soil, and produce other compounds beneficial to microbial growth.

Specific root compounds may be released that have the ability to support and stimulate the growth and metabolic activities of bacteria, including those that can degrade recalcitrant organic compounds. Other root secretions such as root lysates (e.g. phenolic compound) have the capacity to stimulate genes that produce enzymes responsible for microbial metabolism of persistent organic pollutants. Plants may also release surfactants from roots that can reduce interfacial surface tensions of pollutants and increase their aqueous solubility.

AECOM’s screening process for phytoremediation suitability involves assessing plant root bacteria associations through Next Generation Sequencing (NGS) targeting bacteria genes and understanding natural contaminant degrading microfauna in aquifer and vadose zone ecology.

Controlling groundwater contaminant plumes. The use of stands of deep-rooted trees and grasses to control migration of groundwater contaminant plumes is an innovative approach that is becoming increasingly popular. These systems are cost effective with very low operation and maintenance costs.

Special practices have been developed to train roots to extend through a relatively thick vadose zone so that trees can use groundwater from the capillary fringe. The deep-rooted tree stands create a capture zone—a specific thickness of the saturated zone where the groundwater passing beneath the root-zone is used by the trees.


Dissolved contaminants that enter the capture zone can be removed by phytoremediation processes (e.g. microbial degradation in the plant rhizosphere, sorption to roots, plant up-take and detoxification, phytovolatilization, etc.). This approach is useful for many compounds including chlorinated solvents, petroleum hydrocarbons, 1, 4-dioxane, and MTBE. Staggered multiple rows of trees are planted perpendicular to the direction of groundwater flow at the leading edge of the contaminant plume.

The rate of maximal groundwater removal occurs at stand maturity (canopy closure). Plant transpiration results in a significant vertical hydraulic influence, creating an upward driving force to capture groundwater passing through the aquifer. Thicker capture zones are created by increasing the number of rows of trees. This effect is most pronounced for aquifers with lower hydraulic conductivity.

Phytoremediation of PFAS. AECOM is at the forefront of PFAS uptake and transformation work. In May 2017, AECOM developed an unpublished white paper Plant-Assisted Remediation of PFAS which identified various pathways and mechanisms for PFAS uptake and transformation. Later, we provided additional insight into phytoremediation of PFAS through international conference proceedings in 2017, 2018, 2021-2025, and 2026 ([forthcoming](#)). Ongoing research includes mining for novel rhizospheric bacteria in PFAS plumes and role of other natural soil and plant tissue factors in sequestering and altering PFAS.

Key AECOM attributes

- Phytoremediation falls under the NbS umbrella and helps improve native habitat, increase biodiversity and improve upon the human condition.
- AECOM has focused NbS teams in the Great Lakes area, California and the East Coast.
- AECOM phytoremediation services extend from site screening and site suitability, bench-scale, pilot-scale and full-scale applications.



More Information:

Barry J. Harding, CPG, Director Nature Based Solutions, Remediation West | Barry.Harding@aecom.com

Clara Foutch, Project Manager & Ecologist, Remediation West | Clara.Foutch@aecom.com